

What makes KEVZARA different from most other biologics?

KEVZARA targets the interleukin-6 (IL-6) receptor.

Your doctor may prescribe KEVZARA after you've tried a TNFi biologic and it's no longer working well, or it's causing side effects that led you to stop taking the medicine.

▶ **LET'S TAKE A LOOK**

WHAT IS KEVZARA?

KEVZARA[®] (sarilumab) is an injectable prescription medicine called an interleukin-6 (IL-6) receptor blocker. KEVZARA is used to treat adult patients with moderately to severely active rheumatoid arthritis (RA) after at least one other medicine called a disease-modifying antirheumatic drug (DMARD) has been used and did not work well or could not be tolerated.

IMPORTANT SAFETY INFORMATION

KEVZARA can cause serious side effects including:


- **SERIOUS INFECTIONS:** KEVZARA is a medicine that affects your immune system. KEVZARA can lower the ability of your immune system to fight infections. Some people have had serious infections while using KEVZARA, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses that can spread throughout the body. Some people have died from these infections. Your healthcare provider should test you for TB before starting KEVZARA. Your healthcare provider should monitor you closely for signs and symptoms of TB during treatment with KEVZARA.

Please see additional Important Safety Information on the following pages and accompanying full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS** and [Medication Guide](#).

IL-6 and its effects on rheumatoid arthritis (RA).

So what is IL-6? And what does it have to do with your RA?

Interleukin-6, or IL-6, is one of the proteins that's produced by white blood cells. IL-6 plays a major role in the symptoms of RA such as pain, swollen joints, morning stiffness, and fatigue.



When you have RA, IL-6 is produced in excess in inflamed joints and is carried in the blood to other organs. IL-6, along with other proteins (like TNF) and other cells (like T cells), can contribute to the inflammation that may lead to increased disease activity, disability, and joint damage seen in RA.



IL-6

Interleukin 6 (IL-6) is a protein produced by the immune system white cells in your blood. IL-6 is believed to play a role in the symptoms of RA such as pain, swollen joints, morning stiffness, and fatigue.



T cells

A type of white blood cell that, among other things, helps to regulate the immune system. RA is an autoimmune disease where T cells recognize parts of the body as foreign rather than as its own self.



TNF

Tumor Necrosis Factor (TNF) is one of the many proteins produced by the immune system that promote inflammation and may contribute to the joint damage seen in RA.

▶ WHAT ELSE CAN AFFECT YOUR RA?

▶ SO HOW DOES IL-6 CONTRIBUTE TO INFLAMMATION?

IMPORTANT SAFETY INFORMATION (cont'd)

- Before starting KEVZARA (sarilumab), tell your healthcare provider if you
 - think you have an infection or have symptoms of an infection, with or without a fever. Symptoms may include sweats or chills, muscle aches, a cough, shortness of breath, blood in your phlegm, weight loss, warm, red, or painful skin or sores on your body, diarrhea or stomach pain, burning when you urinate or urinating more often than normal, if you feel very tired, or if you are being treated for an infection, get a lot of infections or have repeated infections
 - have diabetes, HIV, or a weakened immune system
 - have TB, or have been in close contact with someone with TB
 - live or have lived, or have traveled to certain parts of the country (such as the Ohio and Mississippi River valleys and the Southwest) where there is an increased chance of getting certain fungal infections (histoplasmosis, coccidioidomycosis, or blastomycosis)
 - have or have had hepatitis
- After starting KEVZARA, call your healthcare provider right away if you have any symptoms of an infection.

Please see additional Important Safety Information on the following pages and accompanying full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS** and [Medication Guide](#).


IMPORTANT SAFETY INFORMATION (cont'd)

- **CHANGES IN CERTAIN LABORATORY TEST RESULTS:** Your healthcare provider should do blood tests before and after starting KEVZARA (sarilumab) to check for low neutrophil (white blood cells that help the body fight off bacterial infections) counts, low platelet (blood cells that help with blood clotting and stop bleeding) counts, and an increase in certain liver function tests. Changes in test results are common with KEVZARA and can be severe. You may also have changes in other laboratory tests, such as your blood cholesterol levels. Your healthcare provider should do blood tests 4 to 8 weeks after starting KEVZARA and then every 6 months during treatment to check for an increase in blood cholesterol levels.
- **TEARS (PERFORATION) OF THE STOMACH OR INTESTINES:** Tell your healthcare provider if you have had a condition known as diverticulitis (inflammation in parts of the large intestine) or ulcers in your stomach or intestines. Some people using KEVZARA had tears in their stomach or intestine. This happens most often in people who also take nonsteroidal anti-inflammatory drugs (NSAIDs), corticosteroids, or methotrexate. Call your healthcare provider right away if you have fever and stomach (abdominal) pain that does not go away.
- **CANCER:** KEVZARA may increase your risk of certain cancers by changing the way your immune system works. Tell your healthcare provider if you have ever had any type of cancer.

Please see additional Important Safety Information on the following pages and accompanying full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS** and [Medication Guide](#).

How IL-6 triggers inflammation.

For a cell to respond to IL-6, IL-6 must connect with an IL-6 receptor.




IL-6R


IL-6 receptor: The IL-6 protein needs to attach to a cell to trigger inflammation. The IL-6 receptor allows the IL-6 protein to connect to the cell.

To help prevent IL-6 from triggering inflammation, these 2 types of receptors need to be blocked.

KEVZARA® is an IL-6 receptor blocker.



The IL-6 receptor can be on the surface of a cell.



Or it can be floating in the bloodstream.

► **HOW CAN KEVZARA HELP?**


IMPORTANT SAFETY INFORMATION (cont'd)

- **SERIOUS ALLERGIC REACTIONS:** Serious allergic reactions can happen with KEVZARA. Get medical attention right away if you have any of the following signs: shortness of breath or trouble breathing; feeling dizzy or faint; swelling of your lips, tongue, or face; moderate or severe stomach (abdominal) pain or vomiting; or chest pain.

Please see additional Important Safety Information on the following pages and accompanying full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS** and [Medication Guide](#).

KEVZARA® interacts with IL-6 in 2 significant ways.

KEVZARA helps to block IL-6 from attaching to receptors that are already on the cell, or while the receptor is in the bloodstream.



In studies of adults with moderate to severe RA who have not been helped by other medicines, **KEVZARA taken with methotrexate has been proven to***:

- Relieve pain in as little as 2 weeks (for some, it may take up to 3 months)
- Reduce tender and swollen joints
- Slow the progression of RA and help protect joints from further damage
- Improve the ability to perform everyday activities

*Individual results may vary.

KEVZARAconnect®

Info and support, when you need it.

We know that starting and staying on a new medication can be difficult. If your doctor thinks KEVZARA is right for you, you'll be able to sign up for the KevzaraConnect® support program, which offers insurance coverage and, if eligible, financial support.

To sign up, visit [KEVZARA.com](https://www.kevzara.com) or call **1-844-KEVZARA** (1-844-538-9272)

IMPORTANT SAFETY INFORMATION (cont'd)

- Do not use KEVZARA if you are allergic to sarilumab or any of the ingredients of KEVZARA.

Please see additional Important Safety Information on the following pages and accompanying full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS** and [Medication Guide](#).

IMPORTANT SAFETY INFORMATION (cont'd)

- Before using KEVZARA, tell your healthcare provider if you
 - have an infection
 - have liver problems
 - have had stomach (abdominal) pain or a condition known as diverticulitis (inflammation in parts of the large intestine) or ulcers in your stomach or intestines
 - recently received or are scheduled to receive a vaccine. People who take KEVZARA should not receive live vaccines
 - plan to have surgery or a medical procedure
 - are pregnant or plan to become pregnant. It is not known if KEVZARA will harm your unborn baby
 - are breastfeeding or plan to breastfeed. Talk to your healthcare provider about the best way to feed your baby if you use KEVZARA. It is not known if KEVZARA passes into your breast milk
 - take any prescription or nonprescription medicines, vitamins, or herbal supplements. It is especially important to tell your healthcare provider if you use
 - any other medicines to treat your RA. Using KEVZARA with these medicines may increase your risk of infection
 - medicines that affect the way certain liver enzymes work. Ask your healthcare provider if you are not sure if your medicine is one of these
- The most common side effects include:
 - injection site redness
 - upper respiratory tract infection
 - urinary tract infection
 - nasal congestion, sore throat, and runny nose

These are not all of the possible side effects of KEVZARA. Tell your doctor about any side effect that bothers you or does not go away. You are encouraged to report side effects of prescription drugs to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

To learn more, talk about KEVZARA with your healthcare provider or pharmacist. The FDA-approved Medication Guide and Prescribing Information can be found at KEVZARA.com or by calling **1-844-KEVZARA** (1-844-538-9272).

Please see full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#).