

KEVZARA®
(sarilumab)

**“I made
my move.”**

SHIRLEY | KEVZARA patient

With KEVZARA, I can get up,
get some exercise, and still
spend time with friends.
This is my moment.

KEVZARA is a prescription medicine
for adult patients with moderately
to severely active rheumatoid arthritis (RA)
after at least one other medicine called a
disease-modifying antirheumatic drug
(DMARD) has been used and did not
work well or could not be tolerated.

Please see Important Safety Information
throughout and full Prescribing Information,
including risk of SERIOUS SIDE EFFECTS
and Medication Guide.

“Starting my day with RA
wasn’t always easy.

That's when I felt the most stiffness and pain. And a rough start usually meant a rough day.

I'd had enough.

I talked with my doctor about trying something different. 

SHIRLEY | KEVZARA patient

IMPORTANT SAFETY INFORMATION

KEVZARA can cause serious side effects including:

- **SERIOUS INFECTIONS:** KEVZARA is a medicine that affects your immune system. KEVZARA can lower the ability of your immune system to fight infections. Some people have had serious infections while using KEVZARA, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses that can spread throughout the body. Some people have died from these infections. Your healthcare provider should test you for TB before starting KEVZARA. Your healthcare provider should monitor you closely for signs and symptoms of TB during treatment with KEVZARA.

- If your current treatment is no longer providing you with adequate relief, talk with your doctor today about taking a different approach with KEVZARA.

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS and Medication Guide.

KEVZARA®
(sarilumab)

How is KEVZARA different?

KEVZARA blocks a protein produced by your white blood cells known as interleukin-6 (IL-6).

Elevated levels of IL-6 are believed to contribute to symptoms of moderate to severe RA, and tend to be higher in the morning.

In a clinical study, KEVZARA with DMARDs was shown to improve symptoms for people with RA who were not experiencing the results they needed with other treatments.

Is it time for a change?

- ▶ Learn more about KEVZARA and how it works differently at KEVZARA.com

IMPORTANT SAFETY INFORMATION (cont'd)

- Before starting KEVZARA, tell your healthcare provider if you
 - think you have an infection or have symptoms of an infection, with or without a fever. Symptoms may include sweats or chills, muscle aches, a cough, shortness of breath, blood in your phlegm, weight loss, warm, red, or painful skin or sores on your body, diarrhea or stomach pain, burning when you urinate or urinating more often than normal, if you feel very tired, or if you are being treated for an infection, get a lot of infections or have repeated infections

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS and Medication Guide.



Different types of biologics target different cells and proteins.

If your current treatment stops working, doesn't work as well as it used to, or if your body can't tolerate it, a different type of biologic may help.

KEVZARA®
(sarilumab)

I'm getting back my day from RA.

Sure, RA affects me, but KEVZARA has helped me start accomplishing ordinary, everyday things again.

So I can get back to the things I love.

SHIRLEY | KEVZARA patient

IMPORTANT SAFETY INFORMATION (cont'd)

- Before starting KEVZARA, tell your healthcare provider if you (cont'd)
 - have diabetes, HIV, or a weakened immune system
 - have TB, or have been in close contact with someone with TB
 - live or have lived, or have traveled to certain parts of the country (such as the Ohio and Mississippi River valleys and the Southwest) where there is an increased chance of getting certain fungal infections (histoplasmosis, coccidioidomycosis, or blastomycosis)
 - have or have had hepatitis
- After starting KEVZARA, call your healthcare provider right away if you have any symptoms of an infection.

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS and Medication Guide.

Why KEVZARA?



KEVZARA comes in a pre-filled pen and pre-filled syringe, both designed with your needs in mind.

Depending on what you and your doctor decide, it can be taken **with or without methotrexate, and is self-injected once every 2 weeks.**

KEVZARA with methotrexate may*:


- Reduce pain after only 2 weeks (although for some, it may take up to 3 months)
- Make joints less sore and swollen
- Slow RA progression and help joints from further damage
- Help improve ability to perform everyday activities

*Individual results may vary.

Both the pen and syringe received an Ease of Use Commendation from the Arthritis Foundation after independent testing by experts and evaluation by patients with arthritis.



KEVZARA®
(sarilumab)



► With the **KevzaraConnect Copay Card**, eligible patients can pay a **\$0 copay per month.***

Maximum copay assistance of \$15,000 per calendar year. Subject to program terms and conditions.

Call **1-844-KEVZARA** (1-844-538-9272) or visit KevzaraConnect.com to learn more and find out if you're eligible.

*Applicable for eligible patients with commercial insurance not funded through a government healthcare program. Subject to an annual maximum copay assistance amount of \$15,000. This program is not valid for prescriptions covered by or submitted for reimbursement under Medicaid, Medicare, VA, DOD, TRICARE, or similar federal or state programs, including any state pharmaceutical assistance program. This program is not valid where prohibited by law, taxed, or restricted. KevzaraConnect reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. See full program terms and conditions at KEVZARA.com.

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS and Medication Guide.

Everyday tips for every day with RA

Morning:

Everyone's different, but most people with RA experience morning stiffness.

To help you rise and shine, try:

- Warming your joints with a heating blanket
 - Stretching in bed
 - Taking a warm shower or bath
- ♥ Morning stretches don't have to be done alone. They can be a way to start off the morning with others.

Noon:

RA may make daily tasks more challenging, but there are ways to make your home work with you.

Go at your own pace and consider:

- Organizing your kitchen so you can easily reach what you use most
 - Replacing heavy shower doors with easy-to-pull curtains
- ♥ Some kitchen appliances (like electric can openers, food processors, and easy-to-grip pots) may help ease the burden of cooking.

Night:

Getting enough rest is key to having a good day, but sleep isn't always easy for people with RA.

Wind down at night by:

- Sticking to a set bedtime
- Listening to soothing music
- Using lighter bedding

- ♥ Writing down your feelings at the end of the day is a good outlet and may help clear your mind.

WHAT IS KEVZARA?

KEVZARA® (sarilumab) is an injectable prescription medicine called an interleukin-6 (IL-6) receptor blocker. KEVZARA is used to treat adult patients with moderately to severely active rheumatoid arthritis (RA) after at least one other medicine called a disease-modifying antirheumatic drug (DMARD) has been used and did not work well or could not be tolerated.

IMPORTANT SAFETY INFORMATION (cont'd)

- **CHANGES IN CERTAIN LABORATORY TEST RESULTS** (cont'd): Changes in test results are common with KEVZARA and can be severe. You may also have changes in other laboratory tests, such as your blood cholesterol levels. Your healthcare provider should do blood tests 4 to 8 weeks after starting KEVZARA and then every 6 months during treatment to check for an increase in blood cholesterol levels.
 - **TEARS (PERFORATION) OF THE STOMACH OR INTESTINES:** Tell your healthcare provider if you have had a condition known as diverticulitis (inflammation in parts of the large intestine) or ulcers in your stomach or intestines. Some people using KEVZARA had tears in their stomach or intestine. This happens most often in people who also take nonsteroidal anti-inflammatory drugs (NSAIDS), corticosteroids, or methotrexate. Call your healthcare provider right away if you have fever and stomach (abdominal) pain that does not go away.
 - **CANCER:** KEVZARA may increase your risk of certain cancers by changing the way your immune system works. Tell your healthcare provider if you have ever had any type of cancer.
 - **SERIOUS ALLERGIC REACTIONS:** Serious allergic reactions can happen with KEVZARA. Get medical attention right away if you have any of the following signs: shortness of breath or trouble breathing; feeling dizzy or faint; swelling of your lips, tongue, or face; moderate or severe stomach (abdominal) pain or vomiting; or chest pain.
 - Do not use KEVZARA if you are allergic to sarilumab or any of the ingredients of KEVZARA.
- Before using KEVZARA, tell your healthcare provider if you
 - have an infection
 - have liver problems
 - have had stomach (abdominal) pain or a condition known as diverticulitis (inflammation in parts of the large intestine) or ulcers in your stomach or intestines
 - recently received or are scheduled to receive a vaccine. People who take KEVZARA should not receive live vaccines
 - plan to have surgery or a medical procedure
 - are pregnant or plan to become pregnant. It is not known if KEVZARA will harm your unborn baby
 - are breastfeeding or plan to breastfeed. Talk to your healthcare provider about the best way to feed your baby if you use KEVZARA. It is not known if KEVZARA passes into your breast milk
 - take any prescription or nonprescription medicines, vitamins, or herbal supplements. It is especially important to tell your healthcare provider if you use
 - any other medicines to treat your RA. Using KEVZARA with these medicines may increase your risk of infection
 - medicines that affect the way certain liver enzymes work. Ask your healthcare provider if you are not sure if your medicine is one of these
- The most common side effects include: injection site redness, upper respiratory tract infection, urinary tract infection, nasal congestion, sore throat, and runny nose

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS and Medication Guide.



Missing out on things you used to find routine?



Don't wait for symptoms to feel unmanageable.

Ask your doctor these quick questions and check out **KEVZARA.com** for more information.

- ▷ What can I do if my current treatment is no longer providing me with the symptom relief I need?
- ▷ Could trying a different kind of treatment help with how I do some everyday activities like running errands, cooking, or going for a walk?
- ▷ How could other treatments, like KEVZARA, be right for me?

IMPORTANT SAFETY INFORMATION (cont'd)

These are not all the possible side effects of KEVZARA. Tell your doctor about any side effect that bothers you or does not go away. You are encouraged to report side effects of prescription drugs to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

To learn more, talk about KEVZARA with your healthcare provider or pharmacist. The FDA-approved Medication Guide and Prescribing Information can be found at **KEVZARA.com** or by calling **1-844-KEVZARA** (1-844-538-9272).

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS and Medication Guide.

SANOFI GENZYME  **REGENERON**