

KEVZARA[®]
(sarilumab) injection
200 mg



Because life
should be a

*Moving
Experience*

Not an actual patient.

KEVZARA is used to treat adults with polymyalgia rheumatica (PMR) after corticosteroids (steroids) have been used and did not work well or when a slow decrease in the dose of corticosteroids (taper) cannot be tolerated.

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS, and Medication Guide at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).

What is KEVZARA?

KEVZARA® (sarilumab) is an injectable prescription medicine called an interleukin-6, or IL-6, receptor blocker.

KEVZARA is the first biologic therapy approved for the treatment of PMR.

KEVZARA is not a steroid.



Not an actual patient.

IMPORTANT SAFETY INFORMATION

KEVZARA can cause serious side effects including:

- **SERIOUS INFECTIONS:** KEVZARA is a medicine that affects your immune system. KEVZARA can lower the ability of your immune system to fight infections. Some people have had serious infections while using KEVZARA, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses that can spread throughout the body. Some people have died from these infections. Your healthcare provider should test you for TB before starting KEVZARA. Your healthcare provider should monitor you closely for signs and symptoms of TB during treatment with KEVZARA.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).

KEVZARA[®]
(sarilumab) injection
200 mg

How does KEVZARA work differently?

The role of IL-6 in PMR inflammation

IL-6 is an immune system protein produced by your white blood cells. When there is too much IL-6 in the body, it can cause the immune system to attack healthy cells.

High levels of IL-6 are believed to play an important role in PMR symptoms, including:



Inflammation



Pain

Treatment of PMR starts with the use of steroids. While steroids may provide symptom relief quickly, some people need to take steroids for a long period of time to keep PMR symptoms under control.



Ask your doctor
about KEVZARA.



Not an actual patient.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):


- Before starting KEVZARA, tell your healthcare provider if you
 - think you have an infection or have signs or symptoms of an infection, with or without a fever such as sweats or chills, muscle aches, a cough, shortness of breath, blood in your phlegm, weight loss, warm, red, or painful skin or sores on your body, diarrhea or stomach pain, burning when you urinate or urinating more often than normal, if you feel very tired, or if you are being treated for an infection, get a lot of infections or have repeated infections.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).

KEVZARA[®]
(sarilumab) injection
200 mg

How KEVZARA works to outsmart inflammation

KEVZARA is designed to block IL-6 effects for people with PMR who want to achieve long-term symptom relief.



KEVZARA works differently **to help control your PMR over time, while allowing you to gradually get off steroids.**

KEVZARA challenges painful inflammation
so you're always one step ahead of it.
Learn more at kevzara.com/pmr-how-it-works.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Before starting KEVZARA, tell your healthcare provider if you (cont'd)
 - have diabetes, HIV, or a weakened immune system.
 - have TB, or have been in close contact with someone with TB.

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS, and Medication Guide at KEVZARA.com/PMR.

KEVZARA[®]
(sarilumab) injection
200 mg

Is KEVZARA right for you?

Sometimes, your PMR symptoms may go away with the help of steroids. When they come back, this is known as a “flare.”

Have you had trouble getting off steroids?

You’re not alone. Four out of ten people living with PMR experience a sudden increase in PMR symptoms within a little after two years of treatment with steroids.



If your current treatment plan isn’t working well,
it might be time for a change.



Talk to your doctor
about KEVZARA.



Not an actual patient.

IMPORTANT SAFETY INFORMATION (cont’d)

KEVZARA can cause serious side effects including (cont’d):

- Before starting KEVZARA, tell your healthcare provider if you (cont’d)
 - live or have lived, or have traveled to certain parts of the country (such as the Ohio and Mississippi River valleys and the Southwest) where there is an increased chance of getting certain fungal infections (histoplasmosis, coccidioidomycosis, or blastomycosis).
 - have or have had hepatitis.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).

KEVZARA[®]
(sarilumab) injection
200 mg

Everyday tips for living with PMR

Healthy lifestyle practices may help you manage your PMR symptoms, so that you can keep up with everyday activities. Below are some tips to help you get started:



Follow a **healthy diet**. Eat plenty of fruits, vegetables, whole grains, and protein.



Get physical. Being inactive may worsen PMR-related pain and stiffness.



Try meditating. Meditation or even simple breathing exercises may help you manage stress and reduce anxiety.



Get enough rest. Sleep is necessary for your body's recovery process.



Use devices designed to make daily tasks easier. These include grocery and rolling laundry carts, reaching aids, and long-handled bath brushes.



Always remember to talk to your doctor about how you are feeling during treatment and before you start or modify any exercise routine.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- After starting KEVZARA, call your healthcare provider right away if you have any symptoms of an infection.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).

KEVZARA[®]
(sarilumab) injection
200 mg

How do I take KEVZARA?

KEVZARA can be taken once every 2 weeks in combination with a dose-lowering course of steroids. Your doctor will guide you on how to gradually lower your dose of steroids.



KEVZARA is given as an injection. KEVZARA is available as a single-use pre-filled syringe or single-use pre-filled pen, designed with your needs in mind.

Learn more at [kevzara.com/pmr-how-to-inject](https://www.kevzara.com/pmr-how-to-inject).



Tips for taking KEVZARA

Sticking to your schedule is important. To help stay on schedule, think about keeping an “injection calendar” and marking the dates and times of your injections so that you don’t forget.

See the detailed Instructions for Use in the **Medication Guide** for instructions about the right way to prepare and inject your KEVZARA at home.

IMPORTANT SAFETY INFORMATION (cont’d)

KEVZARA can cause serious side effects including (cont’d):

- **CHANGES IN CERTAIN LABORATORY TEST RESULTS:** Your healthcare provider should do blood tests before and after starting KEVZARA to check for low neutrophil (white blood cells that help the body fight off bacterial infections) counts, low platelet (blood cells that help with blood clotting and stop bleeding) counts, and an increase in certain liver function tests. Changes in test results are common with KEVZARA and can be severe. You may also have changes in other laboratory tests, such as your blood cholesterol levels. Your healthcare provider should do blood tests 4 to 8 weeks after starting KEVZARA and then every 6 months during treatment to check for an increase in blood cholesterol levels.

Please see Important Safety Information throughout and full Prescribing Information, including risk of **SERIOUS SIDE EFFECTS, and Medication Guide at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).**

KEVZARA[®]
(sarilumab) injection
200 mg

Get the KevzaraConnect® Copay Card



Our copay card may make your copay costs more affordable. With the **KevzaraConnect® Copay Card**, eligible patients may pay as little as **\$0 copay per month.***

Maximum copay assistance of \$15,000 per calendar year. Subject to program terms and conditions.

Call **1-844-KEVZARA** (1-844-538-9272) or learn more and find out if you are eligible at [kevzara.com/pmr-copay](https://www.kevzara.com/pmr-copay).

*Applicable for eligible patients with commercial insurance not funded through a government healthcare program. Subject to an annual maximum copay assistance amount of \$15,000. This program is not valid for prescriptions covered by or submitted for reimbursement under Medicaid, Medicare, VA, DOD, TRICARE, or similar federal or state programs, including any state pharmaceutical assistance program. This program is not valid where prohibited by law, taxed, or restricted. KevzaraConnect® reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. See full program terms and conditions at KEVZARA.com.

Apply for the KevzaraConnect® Copay Card.



Not an actual patient.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- **TEARS (PERFORATION) OF THE STOMACH OR INTESTINES:** Tell your healthcare provider if you have had a condition known as diverticulitis (inflammation in parts of the large intestine) or ulcers in your stomach or intestines. Some people using KEVZARA get tears in their stomach or intestine. This happens most often in people who also take nonsteroidal anti-inflammatory drugs (NSAIDS), corticosteroids, or methotrexate. Call your healthcare provider right away if you have fever and stomach (abdominal) pain that does not go away.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).

KEVZARA®
(sarilumab) injection
200 mg

What are the possible side effects?

KEVZARA can cause serious side effects, including serious allergic reactions. Get medical attention right away if you have any of the following signs of a serious allergic reaction:

- shortness of breath or trouble breathing
- swelling of the lips, tongue, or face
- chest pain
- feeling dizzy or faint
- moderate or severe stomach (abdominal) pain or vomiting

Common side effects of KEVZARA include:

- injection site redness
- injection site itching
- upper respiratory tract infection
- urinary tract infection
- nasal congestion, sore throat, and runny nose

These are not all of the possible side effects of KEVZARA.



Even if they're not listed in this guide, you should always tell your doctor about any changes or side effects you may have.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- **CANCER:** KEVZARA may increase your risk of certain cancers by changing the way your immune system works. Tell your healthcare provider if you have ever had any type of cancer.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).

KEVZARA[®]
(sarilumab) injection
200 mg

Things to know before starting KEVZARA

KEVZARA is a prescription medicine that affects your immune system.
KEVZARA can lower the ability of your immune system to fight infections.

Some people have serious infections while using KEVZARA, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses that can spread throughout the body. Some people have died from these infections.

Your healthcare provider should test you for TB before starting KEVZARA.

- Your healthcare provider should monitor you closely for signs and symptoms of TB during treatment with KEVZARA



You should not start using KEVZARA if you have any kind of infection unless your healthcare provider says it is okay.

Tell your doctor if you think you have an infection or symptoms of an infection, with or without a fever. Signs of an infection may include:

- fever
- sweats or chills
- muscle aches
- cough
- shortness of breath
- blood in your phlegm
- weight loss
- warm, red, or painful skin or sores on your body
- diarrhea or stomach pain
- burning when you urinate or urinating more often than normal
- feeling very tired



Not an actual patient.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- **SERIOUS ALLERGIC REACTIONS:** Serious allergic reactions can happen with KEVZARA. Get medical attention right away if you have any of the following signs: shortness of breath or trouble breathing; feeling dizzy or faint; swelling of your lips, tongue, or face; moderate or severe stomach (abdominal) pain or vomiting; or chest pain.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS, and [Medication Guide](#) at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).**

KEVZARA[®]
(sarilumab) injection
200 mg

What else should you tell your doctor?

Before you start KEVZARA, **make sure your doctor knows about any other infections you currently have or have had in the past**, including if you:

- are being treated for an infection
- get a lot of infections or have infections that keep coming back
- have diabetes, HIV, or a weak immune system. People with these conditions have a higher chance of getting infections
- have TB, or have been in close contact with someone with TB
- live or have lived, or have traveled to certain parts of the country (such as Ohio and Mississippi River valleys and the Southwest) where there is an increased chance of getting certain fungal infections (histoplasmosis, coccidioidomycosis, or blastomycosis). These infections may happen more often or become more severe if you use KEVZARA. Ask your healthcare provider if you do not know if you have lived in an area where these infections are common
- have or have had hepatitis



After starting KEVZARA, call your healthcare provider right away if you have any symptoms of an infection.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Do not use KEVZARA if you are allergic to sarilumab or any of the ingredients of KEVZARA.
- Before using KEVZARA, tell your healthcare provider if you
 - have an infection.
 - have liver problems.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at KEVZARA.com/PMR.

KEVZARA[®]
(sarilumab) injection
200 mg

What does your doctor need to know?

Before using KEVZARA, make sure your doctor knows about any other infections you currently have or have had in the past, including if you:

- have an infection
- have liver problems
- have had stomach (abdominal) pain or been diagnosed with diverticulitis or ulcers in your stomach or intestines
- have recently received or are scheduled to receive a vaccine. People who take KEVZARA should not receive live vaccines
 - All vaccines should be brought up-to-date before starting KEVZARA, unless urgent treatment initiation is required
- plan to have surgery or a medical procedure
- are pregnant or plan to become pregnant. It is not known if KEVZARA will harm your unborn baby
- are breastfeeding or plan to breastfeed. It is not known if KEVZARA passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you use KEVZARA



Tell your doctor about the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Before using KEVZARA, tell your healthcare provider if you (cont'd)
 - have had stomach (abdominal) pain or a condition known as diverticulitis (inflammation in parts of the large intestine) or ulcers in your stomach or intestines.
 - recently received or are scheduled to receive a vaccine. People who take KEVZARA should not receive live vaccines.
 - All vaccines should be brought up-to-date before starting KEVZARA, unless urgent treatment initiation is required.
 - plan to have surgery or a medical procedure.
 - are pregnant or plan to become pregnant. It is not known if KEVZARA will harm your unborn baby.

Please see Important Safety Information throughout and full Prescribing Information, including risk of **SERIOUS SIDE EFFECTS, and Medication Guide at KEVZARA.com/PMR.**

KEVZARA[®]
(sarilumab) injection
200 mg

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Before using KEVZARA, tell your healthcare provider if you (cont'd)
 - are breastfeeding or plan to breastfeed. Talk to your healthcare provider about the best way to feed your baby if you use KEVZARA. It is not known if KEVZARA passes into your breast milk.
 - take prescription or nonprescription medicines, vitamins, or herbal supplements. It is especially important to tell your healthcare provider if you use:
 - any other medicines to treat your RA or PMR. Using KEVZARA with these medicines may increase your risk of infection.
 - medicines that affect the way certain liver enzymes work. Ask your healthcare provider if you are not sure if your medicine is one of these.
- The most common side effects include:
 - injection site redness
 - injection site itching
 - upper respiratory tract infection
 - urinary tract infection
 - nasal congestion, sore throat, and runny nose

These are not all the possible side effects of KEVZARA. Tell your doctor about any side effect that bothers you or does not go away. You are encouraged to report side effects of prescription drugs to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

To learn more, talk about KEVZARA with your healthcare provider or pharmacist. The FDA-approved Medication Guide and Prescribing Information can be found at www.KEVZARA.com or by calling 1-844-KEVZARA.



Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS, and [Medication Guide](#) at KEVZARA.com/PMR.**

KEVZARA[®]
(sarilumab) injection
200 mg

Visit [KEVZARA.com/PMR](https://www.kevzara.com/PMR)
for more information.



Throw PMR out of step with KEVZARA, an IL-6 blocker that challenges painful inflammation, so you're always one step ahead of it.



KEVZARA may help relieve signs and symptoms of PMR when used with a tapering dose of steroids.



KEVZARA is taken once every 2 weeks in combination with a dose-lowering course of steroids.



KEVZARA is self-injected and comes in a single-use pre-filled pen or single-use pre-filled syringe.

Talk to your doctor about how KEVZARA may help.

The most common side effects include:

- injection site redness
- injection site itching
- upper respiratory tract infection
- urinary tract infection
- nasal congestion, sore throat, and runny nose

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS, and Medication Guide at [KEVZARA.com/PMR](https://www.kevzara.com/PMR).

sanofi | **REGENERON**[®]

© 2024 Sanofi and Regeneron Pharmaceuticals, Inc.
All Rights Reserved. MAT-US-2403336-v2.0-07/2024
KEVZARA[®] and KevzaraConnect[®] are registered trademarks of Sanofi Biotechnology.

KEVZARA[®]
(sarilumab) injection
200 mg