

KEVZARA[®]
(sarilumab) injection
200 mg



Because life
should be a

*Moving
Experience*

Not an actual patient.

KEVZARA is used to treat adults with moderately to severely active rheumatoid arthritis (RA) after at least one other medicine called a disease-modifying antirheumatic drug (DMARD) has been used and did not work well or could not be tolerated.

Please see **Important Safety Information** throughout and full **Prescribing Information**, including risk of **SERIOUS SIDE EFFECTS**, and **Medication Guide** at [KEVZARA.com/RA](https://www.kevzara.com/RA).

What is KEVZARA?

KEVZARA® (sarilumab) is an injectable prescription medicine called an interleukin-6, or IL-6, receptor blocker.



KEVZARA with methotrexate has been proven to slow the progression of RA and help protect the joints from further damage.

How is KEVZARA different?




KEVZARA **blocks the effects of a protein** produced by your white blood cells known as **interleukin-6 (IL-6)**.

Elevated levels of IL-6 are believed to contribute to symptoms of moderate-to-severe RA and tend to be higher in the morning.



Ask your doctor
about KEVZARA.



Not an actual patient.

IMPORTANT SAFETY INFORMATION

KEVZARA can cause serious side effects including:


- **SERIOUS INFECTIONS:** KEVZARA is a medicine that affects your immune system. KEVZARA can lower the ability of your immune system to fight infections. Some people have had serious infections while using KEVZARA, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses that can spread throughout the body. Some people have died from these infections. Your healthcare provider should test you for TB before starting KEVZARA. Your healthcare provider should monitor you closely for signs and symptoms of TB during treatment with KEVZARA.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/RA](https://www.kevzara.com/RA).

KEVZARA[®]
(sarilumab) injection
200 mg

How KEVZARA works to outsmart inflammation

KEVZARA is designed to block soluble IL-6 receptors for people with RA to help reduce pain, stiffness, and swelling in the joints.



If your current treatment stops working, doesn't work as well as it used to, or if your body can't tolerate it, KEVZARA may help.

KEVZARA challenges painful inflammation so you're always one step ahead of it.

Learn more at [kevzara.com/ra-how-it-works](https://www.kevzara.com/ra-how-it-works).

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Before starting KEVZARA, tell your healthcare provider if you
 - think you have an infection or have signs or symptoms of an infection, with or without a fever such as sweats or chills, muscle aches, a cough, shortness of breath, blood in your phlegm, weight loss, warm, red, or painful skin or sores on your body, diarrhea or stomach pain, burning when you urinate or urinating more often than normal, if you feel very tired, or if you are being treated for an infection, get a lot of infections or have repeated infections.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/RA](https://www.kevzara.com/RA).

KEVZARA[®]
(sarilumab) injection
200 mg

Why KEVZARA?

KEVZARA, taken with methotrexate or other disease-modifying antirheumatic drugs (DMARDs), may*:



Reduce pain after only 2 weeks (although, for some, it may take up to 3 months).



Make joints less sore and swollen.



Slow the progression of RA and help protect the joints from further damage.

*Individual results may vary.



What this means is that you may be able to perform those everyday activities that matter to you, like making dinner, going to a dance class, or even just taking a nice, long walk. Because life should be a moving experience.

Make sure to talk to your doctor before starting or changing your exercise routine.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Before starting KEVZARA, tell your healthcare provider if you (cont'd)
 - have diabetes, HIV, or a weakened immune system.
 - have TB, or have been in close contact with someone with TB.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/RA](https://www.kevzara.com/RA).

KEVZARA[®]
(sarilumab) injection
200 mg

Everyday tips for managing your RA

Morning



Everyone's different, but most people with RA experience morning stiffness.

To help you rise and shine, try:

- Warming your joints with a heating blanket
- Stretching in bed
- Taking a warm shower or bath

Noon



RA may make daily tasks more challenging, but there are ways to make your space work for you.

Go at your own pace and consider:

- Organizing your kitchen so you can easily reach what you use most
- Replacing heavy shower doors with easy-to-pull curtains

Night



Getting enough rest is key to having a good day, but sleep isn't always easy for people with RA.

Wind down at night by:

- Listening to soothing music
- Using lighter bedding



See more tips on how to manage your RA at [kevzara.com/ra-tips](https://www.kevzara.com/ra-tips).

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Before starting KEVZARA, tell your healthcare provider if you (cont'd)
 - live or have lived, or have traveled to certain parts of the country (such as the Ohio and Mississippi River valleys and the Southwest) where there is an increased chance of getting certain fungal infections (histoplasmosis, coccidioidomycosis, or blastomycosis).
 - have or have had hepatitis.

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS, and Medication Guide at [KEVZARA.com/RA](https://www.kevzara.com/RA).

KEVZARA[®]
(sarilumab) injection
200 mg

How do I take KEVZARA?

KEVZARA comes in a single-use pre-filled pen and single-use pre-filled syringe, both designed with your needs in mind.

Depending on what you and your doctor decide, KEVZARA can be taken with or without methotrexate, and is self-injected once every 2 weeks.

Learn more at kevzara.com/ra-how-to-inject.



Both the pen and syringe received an Ease of Use Commendation from the Arthritis Foundation after independent testing by experts and evaluation by patients with arthritis.



The safety of KEVZARA has been confirmed through multiple studies **over 10 years in RA.**



Not an actual patient.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- After starting KEVZARA, call your healthcare provider right away if you have any symptoms of an infection.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS, and [Medication Guide](#) at KEVZARA.com/RA.**

KEVZARA[®]
(sarilumab) injection
200 mg

Get the KevzaraConnect® Copay Card



Our copay card may make your copay costs more affordable. With the **KevzaraConnect® Copay Card**, eligible patients may pay as little as **\$0 copay per month**.*

Maximum copay assistance of \$15,000 per calendar year. Subject to program terms and conditions.

Call **1-844-KEVZARA** (1-844-538-9272) or learn more and find out if you are eligible at kevzara.com/ra-copay.

*Applicable for eligible patients with commercial insurance not funded through a government healthcare program. Subject to an annual maximum copay assistance amount of \$15,000. This program is not valid for prescriptions covered by or submitted for reimbursement under Medicaid, Medicare, VA, DOD, TRICARE, or similar federal or state programs, including any state pharmaceutical assistance program. This program is not valid where prohibited by law, taxed, or restricted. KevzaraConnect® reserves the right to rescind, revoke, terminate, or amend this offer, eligibility, and terms of use at any time without notice. See full program terms and conditions at KEVZARA.com.

[Apply for the KevzaraConnect® Copay Card.](#)



Not an actual patient.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- **CHANGES IN CERTAIN LABORATORY TEST RESULTS:** Your healthcare provider should do blood tests before and after starting KEVZARA to check for low neutrophil (white blood cells that help the body fight off bacterial infections) counts, low platelet (blood cells that help with blood clotting and stop bleeding) counts, and an increase in certain liver function tests. Changes in test results are common with KEVZARA and can be severe. You may also have changes in other laboratory tests, such as your blood cholesterol levels. Your healthcare provider should do blood tests 4 to 8 weeks after starting KEVZARA and then every 6 months during treatment to check for an increase in blood cholesterol levels.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at KEVZARA.com/RA.

KEVZARA®
(sarilumab) injection
200 mg

What are the possible side effects?

KEVZARA can cause serious side effects, including serious allergic reactions. Get medical attention right away if you have any of the following signs of a serious allergic reaction:

- shortness of breath or trouble breathing
- swelling of the lips, tongue, or face
- chest pain
- feeling dizzy or faint
- moderate or severe stomach (abdominal) pain or vomiting

Common side effects of KEVZARA include:

- injection site redness
- injection site itching
- upper respiratory tract infection
- urinary tract infection
- nasal congestion, sore throat, and runny nose

These are not all of the possible side effects of KEVZARA.



Not an actual patient.

Even if they're not listed in this guide, you should always tell your doctor about any changes or side effects you may have.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- **TEARS (PERFORATION) OF THE STOMACH OR INTESTINES:** Tell your healthcare provider if you have had a condition known as diverticulitis (inflammation in parts of the large intestine) or ulcers in your stomach or intestines. Some people using KEVZARA get tears in their stomach or intestine. This happens most often in people who also take nonsteroidal anti-inflammatory drugs (NSAIDs), corticosteroids, or methotrexate. Call your healthcare provider right away if you have fever and stomach (abdominal) pain that does not go away.

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS, and Medication Guide at [KEVZARA.com/RA](https://www.kevzara.com/RA).

KEVZARA[®]
(sarilumab) injection
200 mg

Things to know before starting KEVZARA

KEVZARA is a prescription medicine that affects your immune system. **KEVZARA can lower the ability of your immune system to fight infections.**

Some people have serious infections while using KEVZARA, including tuberculosis (TB), and infections caused by bacteria, fungi, or viruses that can spread throughout the body. Some people have died from these infections.

Your healthcare provider should test you for TB before starting KEVZARA.

- Your healthcare provider should monitor you closely for signs and symptoms of TB during treatment with KEVZARA



You should not start using KEVZARA if you have any kind of infection unless your healthcare provider says it is okay.

Tell your doctor if you think you have an infection or symptoms of an infection, with or without a fever. Signs of an infection may include:

- fever
- sweats or chills
- muscle aches
- cough
- shortness of breath
- blood in your phlegm
- weight loss
- warm, red, or painful skin or sores on your body
- diarrhea or stomach pain
- burning when you urinate or urinating more often than normal
- feeling very tired

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- **CANCER:** KEVZARA may increase your risk of certain cancers by changing the way your immune system works. Tell your healthcare provider if you have ever had any type of cancer.

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS, and Medication Guide at [KEVZARA.com/RA](https://www.kevzara.com/RA).

KEVZARA[®]
(sarilumab) injection
200 mg

What else should you tell your doctor?

Before you start KEVZARA, **make sure your doctor knows about any other infections you currently have or have had in the past**, including if you:

- are being treated for an infection
- get a lot of infections or have infections that keep coming back
- have diabetes, HIV, or a weak immune system. People with these conditions have a higher chance of getting infections
- have TB, or have been in close contact with someone with TB
- live or have lived, or have traveled to certain parts of the country (such as Ohio and Mississippi River valleys and the Southwest) where there is an increased chance of getting certain fungal infections (histoplasmosis, coccidioidomycosis, or blastomycosis). These infections may happen more often or become more severe if you use KEVZARA. Ask your healthcare provider if you do not know if you have lived in an area where these infections are common
- have or have had hepatitis



Not an actual patient.

After starting KEVZARA, call your healthcare provider right away if you have any symptoms of an infection.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- **SERIOUS ALLERGIC REACTIONS:** Serious allergic reactions can happen with KEVZARA. Get medical attention right away if you have any of the following signs: shortness of breath or trouble breathing; feeling dizzy or faint; swelling of your lips, tongue, or face; moderate or severe stomach (abdominal) pain or vomiting; or chest pain.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at [KEVZARA.com/RA](https://www.kevzara.com/RA).

KEVZARA[®]
(sarilumab) injection
200 mg

What does your doctor need to know?

Before using KEVZARA, talk to your doctor about all of your medical conditions, including if you:

- have an infection
- have liver problems
- have had stomach (abdominal) pain or been diagnosed with diverticulitis or ulcers in your stomach or intestines
- have recently received or are scheduled to receive a vaccine. People who take KEVZARA should not receive live vaccines
 - All vaccines should be brought up-to-date before starting KEVZARA, unless urgent treatment initiation is required.
- plan to have surgery or a medical procedure
- are pregnant or plan to become pregnant. It is not known if KEVZARA will harm your unborn baby
- are breastfeeding or plan to breastfeed. It is not known if KEVZARA passes into your breast milk. Talk to your healthcare provider about the best way to feed your baby if you use KEVZARA



Tell your doctor about the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Do not use KEVZARA if you are allergic to sarilumab or any of the ingredients of KEVZARA.
- Before using KEVZARA, tell your healthcare provider if you
 - have an infection.
 - have liver problems.
 - have had stomach (abdominal) pain or a condition known as diverticulitis (inflammation in parts of the large intestine) or ulcers in your stomach or intestines.
 - recently received or are scheduled to receive a vaccine. People who take KEVZARA should not receive live vaccines.
 - All vaccines should be brought up-to-date before starting KEVZARA, unless urgent treatment initiation is required.
 - plan to have surgery or a medical procedure.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS, and [Medication Guide](#) at [KEVZARA.com/RA](https://www.kevzara.com/RA).**

KEVZARA[®]
(sarilumab) injection
200 mg

IMPORTANT SAFETY INFORMATION (cont'd)

KEVZARA can cause serious side effects including (cont'd):

- Before using KEVZARA, tell your healthcare provider if you (cont'd)
 - are pregnant or plan to become pregnant. It is not known if KEVZARA will harm your unborn baby.
 - are breastfeeding or plan to breastfeed. Talk to your healthcare provider about the best way to feed your baby if you use KEVZARA. It is not known if KEVZARA passes into your breast milk.
 - take prescription or nonprescription medicines, vitamins, or herbal supplements. It is especially important to tell your healthcare provider if you use:
 - any other medicines to treat your RA. Using KEVZARA with these medicines may increase your risk of infection.
 - medicines that affect the way certain liver enzymes work. Ask your healthcare provider if you are not sure if your medicine is one of these.
- The most common side effects include:
 - injection site redness
 - injection site itching
 - upper respiratory tract infection
 - urinary tract infection
 - nasal congestion, sore throat, and runny nose

These are not all the possible side effects of KEVZARA. Tell your doctor about any side effect that bothers you or does not go away. You are encouraged to report side effects of prescription drugs to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

To learn more, talk about KEVZARA with your healthcare provider or pharmacist. The FDA-approved Medication Guide and Prescribing Information can be found at www.KEVZARA.com or by calling 1-844-KEVZARA.



Not an actual patient.

Please see Important Safety Information throughout and full [Prescribing Information](#), including risk of **SERIOUS SIDE EFFECTS**, and [Medication Guide](#) at KEVZARA.com/RA.

KEVZARA[®]
(sarilumab) injection
200 mg

Visit [KEVZARA.com/RA](https://www.kevzara.com/RA)
for more information.

Not an actual patient.

Throw RA out of step with KEVZARA, an IL-6 blocker that challenges painful inflammation, so you're always one step ahead of it.

Taken with methotrexate
or other disease-modifying
antirheumatic drugs (DMARDs):



KEVZARA may* help
slow the progression
of RA.



KEVZARA is taken once
every 2 weeks.



KEVZARA is self-injected
and comes in a single-use
pre-filled pen or single-use
pre-filled syringe.

*Individual results may vary.

Talk to your doctor about how KEVZARA may help.

The most common side effects include:

- injection site redness
- injection site itching
- upper respiratory tract infection
- urinary tract infection
- nasal congestion, sore throat, and runny nose

Please see Important Safety Information throughout and full Prescribing Information, including risk of SERIOUS SIDE EFFECTS, and Medication Guide at [KEVZARA.com/RA](https://www.kevzara.com/RA).

sanofi | **REGENERON**[®]

© 2024 Sanofi and Regeneron Pharmaceuticals, Inc.
All Rights Reserved. MAT-US-2401791-v2.0-07/2024
KEVZARA[®] and KevzaraConnect[®] are registered trademarks of Sanofi Biotechnology.

KEVZARA[®]
(sarilumab) injection
200 mg